

Biochemical Changes Induced by Butachlor in the Liver of Fresh Water Fish *Tor tor* (Ham.)

Vishal Rajput*, Mayuri Pachori, Swati Singh
Department of Biotechnology and Biochemistry
SBSPGI, Balawala,
Dehradun, India
Email:- vsrtech488@gmail.com

Abstract

Freshwater fish Tor tor (Ham.) was exposed to lethal (4.121 ppm) and sub-lethal-I (2.060 ppm) and sub-lethal-II (1.030 ppm) concentration of Butachlor for 72 hrs and 45 days durations. Biochemical changes in liver tissue were analyzed after exposure period. The protein level decreased in lethal and sublethal exposures. Significant decrease in glycogen, slight decrease in protein where as increased cholesterol content has been noted to both concentrations exposure.

Keywords : Protein, Butachlor, Tor tor

Introduction

Man has attempted to increase the world's food production to solve the problem of malnutrition. He achieved this by increased use of fertilizer to nourish the plant and by increased use of pesticides to protect them from pests. Recently a large quantity of pesticides and fertilizers are used to nourish the plants and food production. These chemical have entered into the aquatic system and create pollution, which pose a great threat to aquatic organisms. There are several reports regarding the effects of pesticides (Arunachalam et.al., 1985) on physiology of fish. The pesticides used in pest control programmes also produce many physiological and biochemical changes in freshwater organism particularly the fish (Girija 1984). Although some data available on the effects of different pesticides on the biochemical aspects of fish gill. The

alteration in biochemical contents in different tissues of fish is due to toxic effects of different heavy metals and pesticides have been reported by many workers (Saxena et.al., 1989, Khan et.al., 1992, Virk and Sharma 1999, Rawat et.al., 2002). Results of controlled laboratory exposures of fish to pesticides and related chemical have revealed that liver is often the organ with highest pesticides concentrations (Duke and Wilson 1971). Although more than 900 commercial pesticides are in general use, fewer than 30 have been examined for their adverse effects on fish liver (Pimental et.al., 1971& Gupta 1986).

Tor tor is a very important game fish inhabiting in Lake Bhimtal. Hence the present attempt has been made to study the effect of Butachlor toxicity on the biochemical contents of liver of fresh fish *Tor tor* (Ham).

MATERIAL AND METHODS

Tor tor, were collected from Lake Bhimtal situated at Bhimtal city, district Nainital, (Uttarakhand, India). Fishes were brought to laboratory and kept in aquaria for a week using aged water for acclimatization. During acclimatization they were fed on every day with pieces of live earthworms. The Lc 50 values are determined by following the guideline given by Finney (1971) and Anon (1975). The acclimated fishes were exposed to lethal concentration 4.121 ppm and sub-

lethal-I (2.060 ppm) and sub-lethal-II (1.030 ppm) for 72 hrs for 45 days. Simultaneously a control group of healthy fishes were maintained under identical conditions. The fishes were sacrificed immediately at the end of exposure period and liver was isolated and used to investigate biochemical contents under Butachlor stress. Protein content was estimated by Follin phenol reagent method (Lowry et.al., 1951), Glycogen content was analyzed by using Anthrone reagent method (De zwaan *et.al.*, 1972) and Cholesterol content was analyzed by the method described by Kolmer et.al, (1969).

RESULTS AND DISCUSSION

The results obtained in the present investigation are summarized in Table 1 and Graph 1. Liver is the primary organ for detoxification (Hulterer *et.al.*, 1969) and hence it is expected that toxicant could reach there for detoxification and disposal. This results in structural changes in the liver, the arrangement of hepatic cords leading to the alteration of liver metabolism and its biochemical content. The pollutants acts as one kind of stress and organism respond by developing necessary potential occurring in body give first indication of stress. During stress as organism needs sufficient energy which is supplied from reserve food material *i.e.* protein, glycogen, cholesterol etc. Decrease in the protein content was observed throughout the exposure period. The two sublethal exposure results shows the protein content decreases are depend upon the concentration. There is progressive decrease in the protein content with increase in concentration. The toxicity of Butachlor showed a direct correlation with the concentration and time exposure. Similar observation has been observed by Singh and Bhati (1994).




Fig 1: Comparison of control to lethal, sub-lethal-I and sub-lethal-2 treatment




Fig 2: Percentage of changes reported in protein, glycogen and cholesterol

Decrease in protein may be due to the impairment of protein synthesis or increase in the rate of its degradation to amino acids. This may be fed to TCA cycle through amino transferase probably to cope with high energy demands in order to meet the stress condition. The decrease in proteins



might be due to their utilization in cell repair and tissue organization with the formation of lipoproteins, which are important cellular constituents of cell membranes and cell organelles present in cytoplasm (Harper 1983). The decrease in protein content as a result of toxicity stress has already being reported by Borah and Yadav (1995) and Muley *et. al.*, (2007). Singh (1988) reported reduction in protein content of liver of *Clarias batrachus* in response to Malathion and Y-BHC. Saxena *et.al.*, (1989) attributed the decrease in protein content due to decreased protein synthesizing capacity of liver of *Channa punctatus* exposed to carbaryl and malathion. Jones and Kumar (1996) also observed decline in protein content in liver of *Etrophis maculates* under Ekalux stress. Choudhary and Gaur (2001) observed decline in liver protein of *Cyprinus carpio* under sodium fluoride stress. Shobha *et. al.*, (2007) reported decrease in protein, glycogen and lipid contents in the liver of freshwater fish, *Catla catla* under Cadmium Chloride stress.

Kumar 1996). Decrease in protein content may be due to increased proteolysis (Muley *et. al.*, 2007) or it may be due to metabolic utilization of the ketoacids to glucogenesis pathway for synthesis of glucose (Schmidt Nielson, 1975). The alteration in protein value may also be related to some structural changes in the liver, the arrangement of hepatic words leading to the alteration of liver metabolism.

The decrease in liver protein is also attributed to the inhibition of protein synthesis. Liver glycogen content decreased progressively during exposure period, this may be due to toxic stress. During stress an organism needs sufficient energy which is supplied from reserved glycogen. Glycogen is stored in the organism mainly in the liver and muscles in the form of carbohydrate. It may provide a reserve food for acute demands recurring as a result of transient stress (Love 1980). A fall in the glycogen level clearly indicates its rapid utilization to meet the enhanced energy demands in fish exposed to toxicants through glycolysis or hexose monophosphate pathway. It is assumed that decrease in glycogen content may be due to the inhibition of hormones

Table 1: Biochemical changes induced by Butachlor in the liver of *Tor tor* (HAM).

Parameters	Control	Lethal 4.121 ppm	Change in %	Sub-lethal I 2.060 ppm	Change in %	Sub-lethal II 1.030 ppm	Change in %
Protein	71.78±2.045	59.22**±1.273	-17.5	50.22±2.229	-30.14	55.82±1.097	-22.23
Glycogen	2.166±1.281	0.92**±2.283	-57.16	1.403±1.258	-35.2	1.58±2.008	-26.92
Cholesterol	32.68±1.983	19.30*±2.483	-40.94	21.31±1.829	-34.79	29.91±1.023	-8.33

The values are expressed in mg/100 mg dry weight (mean±SD) [*=P<0.05; **=P<0.01.]

The decrease in protein liver during dimethoate exposure may be due to increased catabolism (Ghousia Begum and Vijaya Raghawan 1995) and decreased anabolism of proteins (Khare and Singh 2002). The loss of protein under Butachlor stress noticed in the present study may be due to the utilization of amino acids in the various catabolic reactions (Jones and

of carbohydrates by the tissue possibly to overcome the pesticides induced stress. James and Sampath (1995) observed decreased liver glycogen in the *Heteropneustes fossilis* (Bloch) under mixtures of copper and ammonia and reported glycogenolysis releasing glucose in to the circulatory system to meet increased energy demand during stress conditions Susan *et. al.*, (1999) observed drastic decreased glycogen content in liver of *Catla catla* under fenvalerate toxicity stress. Rawat



et.al. (2002) have reported continuous decrease in quantity of glycogen in the liver of *Heteropneustes fossilis* exposed to endosulfan. Tiwari and Singh (2009) observed decrease in total protein and glycogen in the liver of *Colisa fasciatus*, exposed to ethanolic extract of Nerium indicum mill latex. Decrease in the glycogen level in liver suggests the possibility of glycogenolysis. A study indicating such depletion in fish models (Mishra and Srivastava 1984) during organophosphorus toxicity offers an excellent support to the decreasing levels of glycogen in the present study.

Cholesterol content was decreased in liver during exposure period. It might be possible that dimethoate causes general damage and structural changes in liver and it leads to effect on capacity of liver to store cholesterol. Khan *et. al.*, (1992) observed significant decrease of cholesterol in liver of Cd treated fish Garra mullya and stated in may be due to general damage in liver. Shakoori *et. al.*, (1996) studied effect of sublethal dose of fenvalerate on the liver of fish Ctenopharyngodon idella and observed decreased level of cholesterol. Virk and Sharma (1999) studied biochemical changes induced by nickel and chromium in the liver of *Cyprinus carpio* and observed significant decline in the cholesterol level of liver and stated this may be due to toxicity stress which suppresses the activity of a number of enzymes responsible for lipid transformation ultimately causing disturbance in lipid metabolism and leads decrease in values of cholesterol. Ascorbic acid content plays an important role in detoxification of the foreign bodies or toxicants in metabolic process. The main process of detoxification takes place in liver and is also the main site where ascorbic acids are synthesized. Due to pesticidal stress significant increase in ascorbic content was observed in liver of

Tor tor during present study. Bhusari (1985) observed increased amount of ascorbic acid in liver of *Barbus ticto* during endosulfan, ekalux and sevmol exposure. Davane (1991) observed increased level of liver ascorbic acid in *T. sandkhol* to dimethoate, thiodon and carbaryl. The liver is the main site of detoxification and having large amount of ascorbic acid. This type of observation is also made by Somasundaram *et. al.*, (1978). The ascorbic acid plays a role directly related to homeostatic mechanism and is essential for wound healing and regeneration. Gould (1963) Shah *et. al.*, (1971) several investigations have reported protective effects of ascorbic acid against the toxicity of various environmental chemicals Padhi and Pathaik (1978). Due to this factor ascorbic acid content must have been increased in liver of *Tor tor* under Butachlor toxicity stress.

Acknowledgement

Authors are thankful to the Sardar Bhagwan Singh (PG) Institute of Biomedical Sciences and Reaserch, Balawala, Dehradun for providing laboratory and instrumentation facilities.

REFERENCES

1. Anon, J. (1975). Recommendations of the committee on methods for toxicity test with fish, macro-invertebrates and amphibians EPA, Oregon. 61pp.
2. Arunachalam, S., Palanichamy, S., Balasubramanian, M.P. (1985): Sublethal effects of carbaryl on food utilization and oxygen consumption in the air breathing fish *Channa punctatus* (Bloch). Journal Environmental Biology. 6, (4), 279-286.
3. Bhusari, N.B. (1985). Studies on the effect of pesticides on freshwater fish of Marathwada Region Ph.D. thesis Dr. BAMU, Aurangabad (MS) India.
4. Borah, S. Yadav, R.N.S. (1995). Alterations in the protein free amino acid, nucleic acids and carbohydrate contents of muscle and gill in rogor exposed freshwater fish *Heteropneustes fossilis*. Pollution Research. 14, (1), 99-103.
5. Choudhary, A. Gaur, S. (2001). Effect of sodium fluoride on the muscle and liver of a freshwater fish, *Cyprinus carpio*. Journal Aqua. Biol. 16, (2), 67-68.
6. Dang, J.N., Kumar, S.G. (1996). Effect of Ekalux on biochemical parameters in the freshwater fish *Etroplus maculatus*. Journal of Ecotoxicology and Environmental Monitoring. 6, (1), 65-67.

7. Davane, P.M. (1991). Impacts of assessment of pesticides in aquatic animals, special reference with fishes Ph.D. thesis Dr. BAMU, Aurangabad (MS) India.
8. Duke, T.W., Wilson, A.J. (1971). Chlorinated hydrocarbons in liver of fishes from the northeastern Pacific Ocean. *Pestic. Monit. J. Sci.*, 228-232.
9. Finney, D.J. (1971). Probit analysis London Cambridge university press. pp-333.
10. Ghousia, B., Vijayaraghavan, S. (1995). In vivo toxicity of dimethoate on proteins and transaminases, in the liver tissue of fresh water fish, *Clarias batrachus* (Linn.) *Bulletin of Environmental Contamination Toxicology*. 54, 370-375.
11. Girija, M. (1984). Effect of Heptachlor and metasytox on fish *Tilapia mossambica* (Petero). M.Phil Dissertation Sri Venkateswara University, Tirupati, India.
12. Gould, B.S. (1963). Collagen formation and fibrogenesis with special reference to the role of ascorbic acid Gut. *Review of Cytology*. 15, 301-363.
13. Gupta, P.K. (1986). Pesticides in Indian environment interprint, New Delhi.
14. Harper, A.H. (1983). *Review of Bio Chemistry* 20th ed. Lange medical Publication Co. California, 1012 pp.
15. Hulterer, F., Klion, F.M., Wengraf, A., Schaffner, F., Poper, H., (1969). Hepatocellular adaptation and injury structural and biochemical changes following dieldrin and methyl butter yellow. *Lab. Invest.* 20, 455-464.
16. James, R., Sampath, K. (1995). Sublethal effects of mixtures of copper and ammonia on selected biochemical and physiological parameters in the catfish *Heteropneustes fossilis* (Bloch). *Bulletin of Environmental Contamination Toxicology*. 55, 187-194.
17. Khan, E.A., Sinha, M.P., Saxena, N. Mehrotra, P.N. (1992). Biochemical effect of cadmium toxicity on a Hill stress Teleost *Garra mullya* (sykes) during Egg maturation. *Cholesterol and Glycogen, Pollution Research*. 11, (3), 163-167.
18. Khare, A., Singh, S. (2002). Impact of malathion on protein content in the freshwater fish *Clarias batrachus*. *Journal of Ecotoxicology and Environmental Monitoring*. 12, (2), 129-132.
19. Kolmer, J.A., Spaulding, E.H., Robinson, H.W. (1969). *Approved Laboratory Technique*, scientific book agencies, Calcutta (India).
20. Love, R.M. (1980). *The chemical biology of fishes* academic press, New York. 943 pp.
21. Lowry, O.H., Rosenbrough, N.J., Forr, A.L., Randal, R.J., (1951). Protein measurement with Folin Phenol Reagent. *Journal Biol. Chem.* 193, 265-275.
22. Mishra, J., Srivatava, A.K. (1984). Effects of chlordane on the blood and tissue chemistry of a teleost fish, *Heteropneustes fossilis*. *Cell and Molecular Biology*, 30, 519-523.
23. Muley, D.V., Karanjkar, D.M., Maske, S.V. (2007). Impact of industrial effluents on the biochemical composition of freshwater fish *Labeo rohita*. *Journal Environmental Biology*. 28, (2), 245-249.
24. Padhi, S.M., Patnaik, B.K., (1978). Changes in ascorbic acid content kidney following partial hepatectomy in the garden lizard *Calotes versicolor* (Daudin). *Indian Journal Experimental Biology*. 16, 1499.
25. Pimentel, D. (1971). *Ecological effects of pesticides on nontarget species*. Exec of president off sci. technical, U.S. Government printing off. Washington, D.C.
26. Radhaiah, V., Girija, M., Raok, J., (1987). Changes in selected biochemical parameters in the kidney and blood of the fish *Tilapia mossambica* (peters) exposed to heptachlor. *Bulletin of Environmental Contamination Toxicology*. 39, 1006-1011.
27. Rawat, D.K., Bais, V.S., Agrawal, N.C. (2002). A correlative study on liver glycogen and endosulfan toxicity in *Heteropneustes fossilis* (Bioch). *Journal Environmental Biology*. 23, (2), 205-207.
28. Roe, J.H. (1967). *Methods of Biochemical analysis*, Vol. 5: (Ed. By Glick Inter. science, New York), 44 pp.
29. Saxena, P.K., Singh, V.P., Kondal, J.K., Soni, G.L. (1989). Effect of some pesticides on invitro lipid and protein synthesis by the liver of the freshwater teleost *Channa punctatus* (Bloch.). *Environmental Pollution*. 58, 273-276.
30. Schmidt-Nielson, B. (1989). Comparative Physiology of cellular ion and volume regulation. *Journal of Experimental Zoology*. 194, 107-220
31. Shah, R.K., Hiradhar, P.K., Magon, D.K. (1971). Ascorbic acid in the normal and regenerating tail of the house lizard *Hemidactylus flaviviridis*. *Journal Embryology and Experimental Morphology*. 26, 285.
32. Shakoori, A.R., Mughal, A.L., Iqbal, M.J. (1996). Effects of sublethal doses of fenvalerate (A. synthetic pyrethroid, Aadministered continuously for four weeks) on the blood, liver and muscle of a freshwater fish *Ctenopharyngodon idella*. *Bulletin of Environmental Contamination Toxicology*. 57, 487-494.
33. Sing, T.P. (1988). Impact of agricultural pesticides on the enclocrinology of some freshwater fishes. Final report USDA, ICAR New Delhi. India.
34. Singh, S., Bhati, D.P.S. (1994). Evaluation of liver protein due to stress under 2,4-D. Intoxication in *Channa punctatus* (Bioch). *Bulletin of Environmental Contamination Toxicology*. 53, 149-152.
35. Somasundaram, K., Damodaran, V.N., Venkatasubramanian, T.A. (1975). Biochemical changes following absorption of dieldrin through skin. *Indian Journal Experimental Biology*. 16, (9), 1003-1004.
36. Susan, T.A., Veeraiah, K. Tilak, K.S. (1999). Biochemical and enzymatic changes in the tissues of *Catla catla* exposed to the pyrethroid fenvalerate. *Journal Ecobiology*. 11, (2), 109-116.
37. Virk, S., Sharma, R.C. (1999). Biochemical changes induced by Nickel and Chromium in the liver of *Cyprinus carpio* (L). *Pollution Research*. 18, (3), 217-222.
38. Zwaan, D.C.A., Zandee, D.I. (1971). Glycogen estimation with Anthrone reagent comp. *Biochem. physiol.* 43B, 53-55.